arXiv:1403.4447v1 [math.NT] 18 Mar 2014

A NOTE ON q-ANALOGUE OF BOOLE POLYNOMIALS

DAE SAN KIM, TAEKYUN KIM, JONG JIN SEO

Abstract. In this paper, we consider the q-extensions of Boole polynomials. From those polynomials, we derive some new and interesting properties and identities related to special polynomials.

1. Introduction

Let p be a fixed odd prime number. Throughout this paper, $\mathbb{Z}_p, \mathbb{Q}_p$ and $\mathbb{C}_p$ will denote the ring of p-adic integers, the field of p-adic numbers and the completion of algebraic closure of $\mathbb{Q}_p$. The p-adic norm $|\cdot|_p$ is normalized as $|p|_p = 1/p$. The space of continuous functions on $\mathbb{Z}_p$ is denoted by $C(\mathbb{Z}_p)$. Let q be an indeterminate in $\mathbb{C}_p$ with $|1-q|_p < p^{-1/p-1}$. The q-number of x is defined by $[x]_q = \frac{1-q^x}{1-q}$. Note that $\lim_{q\to 1}[x]_q = x$. For $f \in C(\mathbb{Z}_p)$, the fermionic p-adic q-integral on $\mathbb{Z}_p$ is defined by Kim to be

$$\begin{aligned} I_{-q}(f) &= \int_{\mathbb{Z}_p} f(x)d\mu_{-q}(x) = \lim_{N\to\infty}\frac{1}{[p^N]_{-q}}\sum_{x=0}^{p^N-1} f(x)(-1)^x, \\ &\text{where } [x]_{-q} = \frac{1-(-q)^x}{1+q} \text{ (see [1 − 9])}. \end{aligned} \tag{1.1}$$

From (1.1), we note that

$$\begin{aligned} q^n I_{-q}(f_n) + (-1)^{n-1} I_{-q}(f) &= [2]_q \sum_{l=0}^{n-1}(-1)^{n-1-l} q^l f(l), \\ &\text{where } f_n(x) = f(x+n), (n \geq 1) \text{ (see [4])}. \end{aligned} \tag{1.2}$$

In particular, for n=1,

$$qI_{-q}(f_1) + I_{-q}(f) = [2]_q f(0). \tag{1.3}$$

Key words and phrases. q-Boole number, q-Boole polynomial, q-Euler number, q-Euler polynomial .

.

As is well known, the Boole polynomials are defined by the generating function to be

$$\sum_{n=0}^{\infty} Bl_n(x|\lambda)\frac{t^n}{n!} = \frac{1}{1+(1+t)^{\lambda}}(1+t)^x, \text{ (see [2, 11])}. \tag{1.4}$$

When $\lambda = 1, 2Bl_n(x|1) = Ch_n(x)$ are Changhee polynomials which are defined by

$$\frac{2}{t+2}(1+t)^x = \sum_{n=0}^{\infty} Ch_n(x)\frac{t^n}{n!} \text{ (see [2])}. \tag{1.5}$$

The Euler polynomials of order α are defined by the generating function to be

$$\left(\frac{2}{e^t+1}\right)^{\alpha} e^{xt} = \sum_{n=0}^{\infty} E_n^{(\alpha)}(x)\frac{t^n}{n!}, \text{ (see [2, 11])}. \tag{1.6}$$

When $x = 0, E_n^{(\alpha)} = E_n^{(\alpha)}(0)$ are called the Euler numbers of order α.
In particular, for $\alpha = 1, E_n(x) = E_n^{(1)}(x)$ are called the ordinary Euler polynomials.
The Stirling number of the first kind is given by the generating function to be

$$\log(1+t)^m = m!\sum_{l=m}^{\infty} S_1(l,m)\frac{t^l}{l!}, (m \geq 0), \tag{1.7}$$

and the Stirling number of the second kind is defined by the generating function to be

$$(e^t-1)^m = m!\sum_{l=m}^{\infty} S_2(l,m)\frac{t^l}{l!}, \text{ (see [11])}. \tag{1.8}$$

In this paper, we consider the q-extensions of Boole polynomials. From those polynomials, we derive new and interesting properties and identities related to special polynomials.

2. q-analogue of Boole polynomials

In this section, we assume that $t \in \mathbb{C}_p$ with $|t|_p < p^{\frac{-1}{p-1}}$ and $\lambda \in \mathbb{Z}_p$ with $\lambda \neq 0$. From (1.3), we note that

$$\begin{aligned}\int_{\mathbb{Z}_p} (1+t)^{x+\lambda y} d\mu_{-q}(y) &= \frac{1+q}{1+q(1+t)^{\lambda}}(1+t)^x \\ &= \sum_{n=0}^{\infty} [2]_q Bl_{n,q}(x|\lambda)\frac{t^n}{n!},\end{aligned} \tag{2.1}$$

where $Bl_{n,q}(x|\lambda)$ are the q-Boole polynomials which are defined by

$$\frac{1}{1+q(1+t)^{\lambda}}(1+t)^{x}=\sum_{n=0}^{\infty}Bl_{n,q}(x|\lambda)\frac{t^{n}}{n!}. \tag{2.2}$$

From (2.1), we can derive the following equation :

$$\int_{\mathbb{Z}_p}\binom{x+\lambda y}{n}d\mu_{-q}(y)=\frac{[2]_q}{n!}Bl_{n,q}(x|\lambda). \tag{2.3}$$

When $x=0, Bl_{n,q}(\lambda)=Bl_{n,q}(0|\lambda)$ are called the q-Boole numbers.
Now, we observe that

$$\begin{aligned}(1+t)^{x+\lambda y}&=e^{(x+\lambda y)\log(1+t)}\\&=\sum_{m=0}^{\infty}\frac{(x+\lambda y)^{m}}{m!}\big(\log(1+t)\big)^{m}\\&=\sum_{m=0}^{\infty}\frac{(x+\lambda y)^{m}}{m!}m!\sum_{n=m}^{\infty}S_1(n,m)\frac{t^{n}}{n!}\\&=\sum_{n=0}^{\infty}\left\{\sum_{m=0}^{n}(x+\lambda y)^{m}S_1(n,m)\right\}\frac{t^{n}}{n!}.\end{aligned} \tag{2.4}$$

The q-Euler polynomials are defined by the generating function to be

$$\frac{[2]_q}{qe^{t}+1}e^{xt}=\sum_{n=0}^{\infty}E_{n,q}(x)\frac{t^{n}}{n!}. \tag{2.5}$$

Note that $\lim_{q\to 1}E_{n,q}(x)=E_n(x)$.
When $x=0, E_{n,q}=E_{n,q}(0)$ are called the q-Euler numbers. By (1.3), we easily get

$$\begin{aligned}\int_{\mathbb{Z}_p}e^{(x+y)t}d\mu_{-q}(y)&=\frac{[2]_q}{qe^{t}+1}e^{xt}\\&=\sum_{n=0}^{\infty}E_{n,q}(x)\frac{t^{n}}{n!}.\end{aligned} \tag{2.6}$$

Thus, by (2.6), we get

$$\int_{\mathbb{Z}_p}(x+y)^{n}d\mu_{-q}(y)=E_{n,q}(x),(n\geq 0). \tag{2.7}$$

From (2.1), (2.4) and (2.7), we have

$$\begin{aligned}\int_{\mathbb{Z}_p}(1+t)^{x+\lambda y}d\mu_{-q}(y) &= \sum_{n=0}^{\infty}\left\{\sum_{m=0}^{n}\int_{\mathbb{Z}_p}(x+\lambda y)^m d\mu_{-q}(y)S_1(n,m)\right\}\frac{t^n}{n!}\\ &= \sum_{n=0}^{\infty}\left\{\sum_{m=0}^{n}\lambda^m E_{m,q}\left(\frac{x}{\lambda}\right)S_1(n,m)\right\}\frac{t^n}{n!}.\end{aligned} \tag{2.8}$$

Therefore, by (2.1), (2.3) and (2.8), we obtain the following theorem.

Theorem 2.1. *For $n \geq 0$, we have*

$$Bl_{n,q}(x|\lambda) = \frac{1}{[2]_q}\sum_{m=0}^{n}\lambda^m E_{m,q}\left(\frac{x}{\lambda}\right)S_1(n,m),$$

and

$$\int_{\mathbb{Z}_p}\binom{x+\lambda y}{n}d\mu_{-q}(y) = \frac{[2]_q}{n!}Bl_{n,q}(x|\lambda).$$

From (2.3), we note that

$$Bl_{n,q}(x|\lambda) = \frac{1}{[2]_q}\int_{\mathbb{Z}_p}(x+\lambda y)_n d\mu_{-q}(y).$$

When $\lambda = 1$, we have

$$Bl_{n,q}(x|1) = \frac{1}{[2]_q}\int_{\mathbb{Z}_p}(x+y)_n d\mu_{-q}(y). \tag{2.9}$$

As is known, q-Changhee polynomials are defined by the generating function to be

$$\frac{[2]_q}{[2]_q+qt}(1+t)^x = \sum_{n=0}^{\infty}Ch_{n,q}(x)\frac{t^n}{n!}. \tag{2.10}$$

Thus, by (2.10), we get

$$\int_{\mathbb{Z}_p}(1+t)^{x+y}d\mu_{-q}(y) = \frac{[2]_q}{[2]_q+qt}(1+t)^x = \sum_{n=0}^{\infty}Ch_{n,q}(x)\frac{t^n}{n!}. \tag{2.11}$$

From (2.11), we have

$$\begin{aligned}&\int_{\mathbb{Z}_p}(x+y)_n d\mu_{-q}(y) = Ch_{n,q}(x),\\ &\text{where}(x)_n = x(x-1)\cdots(x-n+1).\end{aligned} \tag{2.12}$$

By (2.9) and (2.12), we get

$$Bl_{n,q}(x|1) = \frac{1}{[2]_q}Ch_{n,q}(x). \tag{2.13}$$

By replacing t by e^t-1 in (2.2), we see that

$$\begin{aligned}\frac{[2]_q}{qe^{\lambda t}+1}e^{xt} &= [2]_q\sum_{n=0}^{\infty}Bl_{n,q}(x|\lambda)\frac{1}{n!}(e^t-1)^n \\ &= [2]_q\sum_{n=0}^{\infty}Bl_{n,q}(x|\lambda)\sum_{m=n}^{\infty}S_2(m,n)\frac{t^m}{m!} \\ &= \sum_{m=0}^{\infty}\sum_{n=0}^{m}[2]_qBl_{n,q}(x|\lambda)S_2(m,n)\frac{t^m}{m!},\end{aligned} \tag{2.14}$$

and

$$\begin{aligned}\frac{[2]_q}{qe^{\lambda t}+1}e^{xt} &= \frac{[2]_q}{qe^{\lambda t}+1}e^{(\frac{x}{\lambda})\lambda t} \\ &= \sum_{m=0}^{\infty}E_{m,q}\left(\frac{x}{\lambda}\right)\lambda^m\frac{t^m}{m!}.\end{aligned} \tag{2.15}$$

Therefore, by (2.14) and (2.15), we obtain the following theorem.

Theorem 2.2. *For $m\geq 0$, we have*

$$\sum_{n=0}^{m}Bl_{n,q}(x|\lambda)S_2(m,n)=\frac{1}{[2]_q}E_{m,q}\left(\frac{x}{\lambda}\right)\lambda^m.$$

Let us define the q-Boole numbers of the first kind with order $k(\in\mathbb{N})$ as follows :

$$[2]_q^kBl_{n,q}^{(k)}(\lambda)=\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}(\lambda(x_1+\cdots+x_k))_nd\mu_{-q}(x_1)\cdots d\mu_{-q}(x_k),(n\geq 0). \tag{2.16}$$

Thus, by (2.16), we see that

$$\begin{aligned}[2]_q^k\sum_{n=0}^{\infty}Bl_{n,q}^{(k)}(\lambda)\frac{t^n}{n!} &= \int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}\sum_{n=0}^{\infty}\binom{\lambda(x_1+\cdots+x_k)}{n}t^nd\mu_{-q}(x_1)\cdots d\mu_{-q}(x_k) \\ &= \int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}(1+t)^{\lambda(x_1+\cdots+x_k)}d\mu_{-q}(x_1)\cdots d\mu_{-q}(x_k) \\ &= \left(\frac{1+q}{1+q(1+t)^{\lambda}}\right)^k \\ &= [2]_q^k\sum_{n=0}^{\infty}\left(\sum_{l_1+\cdots+l_k=n}\binom{n}{l_1,\cdots,l_k}Bl_{l_1,q}\cdots Bl_{l_k,q}\right)\frac{t^n}{n!}.\end{aligned} \tag{2.17}$$

Therefore, by (2.17), we obtain the following corollary.

Corollary 2.3. *For $n \geq 0$, we have*

$$Bl_{n,q}^{(k)} = \sum_{l_1+\cdots+l_k=n} \binom{n}{l_1,\cdots,l_k} Bl_{l_{1,q}} \cdots Bl_{l_{k,q}}.$$

The q-Euler polynomials of order k are defined by the generating function to be

$$\begin{aligned} &\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} e^{(x_1+\cdots+x_k+x)t} d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k) \\ &= \left(\frac{[2]_q}{qe^t+1}\right)^k e^{xt} = \sum_{n=0}^{\infty} E_{n,q}^{(k)}(x) \frac{t^n}{n!}. \end{aligned} \tag{2.18}$$

Thus, by (2.18), we get

$$\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} (x_1+\cdots+x_k+x)^n d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k) = E_{n,q}^{(k)}(x).$$

When $x = 0, E_{n,q}^{(k)} = E_{n,q}^{(k)}(0)$ are called the q-Euler numbers of order k. From (2.16), we note that

$$\begin{aligned} [2]_q^k Bl_{n,q}^{(k)}(\lambda) &= \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} (\lambda(x_1+\cdots+x_k))_n d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k) \\ &= \sum_{l=0}^{n} S_1(n,l) \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} \lambda^l (x_1+\cdots+x_k)^l d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k) \\ &= \sum_{l=0}^{n} S_1(n,l) \lambda^l E_{l,q}^{(k)}. \end{aligned} \tag{2.19}$$

Therefore, by (2.19), we obtain the following theorem.

Theorem 2.4. *For $n \geq 0$, we have*

$$Bl_{n,q}^{(k)}(\lambda) = \frac{1}{[2]_q^k} \sum_{l=0}^{n} S_1(n,l) \lambda^l E_{l,q}^{(k)}.$$

By replacing t by $e^t - 1$ in (2.17), we get

$$\begin{aligned} [2]_q^k \sum_{n=0}^{\infty} Bl_{n,q}^{(k)}(\lambda) \frac{1}{n!} (e^t-1)^n &= \left(\frac{[2]_q}{qe^{\lambda t}+1}\right)^k \\ &= \sum_{m=0}^{\infty} E_{m,q}^{(k)} \lambda^m \frac{t^m}{m!}, \end{aligned} \tag{2.20}$$

and

$$\begin{aligned}[2]_q^k \sum_{n=0}^{\infty} Bl_{n,q}^{(k)}(\lambda)\frac{1}{n!}(e^t-1)^n &= [2]_q^k \sum_{n=0}^{\infty} Bl_{n,q}^{(k)}(\lambda) \sum_{m=n}^{\infty} S_2(m,n)\frac{t^m}{m!} \\ &= [2]_q^k \sum_{m=0}^{\infty} \left\{ \sum_{n=0}^{m} Bl_{n,q}^{(k)}(\lambda) S_2(m,n) \right\} \frac{t^m}{m!}.\end{aligned} \tag{2.21}$$

Therefore, by (2.20) and (2.21), we obtain the following theorem.

Theorem 2.5. *For $m \geq 0$, we have*

$$\sum_{n=0}^{m} Bl_{n,q}^{(k)}(\lambda) S_2(m,n) = \frac{1}{[2]_q^k} E_{m,q}^{(k)} \lambda^m.$$

Let us define the higher-order q-Boole polynomials of the first kind as follows :

$$[2]_q^k Bl_{n,q}^{(k)}(x|\lambda) = \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} (\lambda x_1 + \cdots + \lambda x_k + x)_n d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k),$$

where $n \geq 0$ and $k \in \mathbb{N}$. (2.22)

From (2.22), we can derive the generating function of the higher-order q-Boole polynomials of the first kind as follows :

$$\begin{aligned}[2]_q^k \sum_{n=0}^{\infty} Bl_{n,q}^{(k)}(x|\lambda)\frac{t^n}{n!} &= \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} (1+t)^{\lambda x_1 + \cdots + \lambda x_k + x} d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k) \\ &= \left(\frac{[2]_q}{1+q(1+t)^{\lambda}} \right)^k (1+t)^x\end{aligned} \tag{2.23}$$

By (2.17), we easily get

$$\begin{aligned}\left(\frac{[2]_q}{1+q(1+t)^{\lambda}} \right)^k (1+t)^x &= [2]_q^k \left(\sum_{l=0}^{\infty} Bl_{l,q}^{(k)}(\lambda)\frac{t^l}{l!} \right) \left(\sum_{m=0}^{\infty} m! \binom{x}{m} \frac{t^m}{m!} \right) \\ &= [2]_q^k \sum_{n=0}^{\infty} \left(\sum_{m=0}^{n} m! \binom{x}{m} \frac{n!}{m!(n-m)!} Bl_{n-m,q}^{(k)}(\lambda) \right) \frac{t^n}{n!} \\ &= [2]_q^k \sum_{n=0}^{\infty} \left(\sum_{m=0}^{n} m! \binom{x}{m} \binom{n}{m} Bl_{n-m,q}^{(k)}(\lambda) \right) \frac{t^n}{n!}.\end{aligned} \tag{2.24}$$

Therefore, by (2.23) and (2.24), we obtain the following theorem.

Theorem 2.6. *For $n \geq 0$, we have*

$$Bl_{n,q}^{(k)}(x|\lambda) = \sum_{m=0}^{n} \binom{n}{m} Bl_{n-m,q}^{(k)}(\lambda)(x)_m.$$

Replacing t by $e^t - 1$ in (2.23), we have

$$\begin{aligned} [2]_q^k \sum_{n=0}^{\infty} Bl_{n,q}^{(k)}(x|\lambda) \frac{(e^t-1)^n}{n!} &= \left(\frac{[2]_q}{1+qe^{\lambda t}} \right)^k e^{xt} \\ &= \sum_{m=0}^{\infty} E_{m,q}^{(k)}\left(\frac{x}{\lambda}\right) \lambda^m \frac{t^m}{m!}, \end{aligned} \tag{2.25}$$

and

$$[2]_q^k \sum_{n=0}^{\infty} Bl_{n,q}^{(k)}(x|\lambda) \frac{(e^t-1)^n}{n!} = [2]_q^k \sum_{m=0}^{\infty} \left(\sum_{n=0}^{m} Bl_{n,q}^{(k)}(x|\lambda) S_2(m,n) \right) \frac{t^m}{m!}. \tag{2.26}$$

Thus, from (2.25) and (2.26), we have the following theorem.

Theorem 2.7. *For $m \geq 0$ and $k \in \mathbb{N}$, we have*

$$\sum_{n=0}^{m} Bl_{n,q}^{(k)}(x|\lambda) S_2(m,n) = \frac{1}{[2]_q^k} \lambda^m E_{m,q}^{(k)}\left(\frac{x}{\lambda}\right).$$

From (2.22), we have

$$\begin{aligned} [2]_q^k Bl_{n,q}^{(k)}(x|\lambda) &= \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} (\lambda x_1 + \cdots + \lambda x_k + x)_n d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k) \\ &= \sum_{l=0}^{n} S_1(n,l) \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} (\lambda x_1 + \cdots + \lambda x_k + x)^l d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k) \\ &= \sum_{l=0}^{n} S_1(n,l) \lambda^l E_{l,q}^{(k)}\left(\frac{x}{\lambda}\right). \end{aligned} \tag{2.27}$$

Therefore, by (2.27), we obtain the following theorem.

Theorem 2.8. *For $n \geq 0$, $k \in \mathbb{N}$, we have*

$$Bl_{n,q}^{(k)}(x|\lambda) = \frac{1}{[2]_q^k} \sum_{l=0}^{n} S_1(n,l) \lambda^l E_{l,q}^{(k)}\left(\frac{x}{\lambda}\right).$$

Now, we consider the q-analogue of Boole polynomials of the second kind as follows :

$$\widehat{Bl}_{n,q}(x|\lambda) = \frac{1}{[2]_q} \int_{\mathbb{Z}_p} (-\lambda y + x)_n d\mu_{-q}(y), (n \geq 0). \tag{2.28}$$

Thus, by (2.28), we get

$$\begin{aligned}\widehat{Bl}_{n,q}(x|\lambda) &= \frac{1}{[2]_q}\sum_{l=0}^{n} S_1(n,l)(-1)^l\lambda^l \int_{\mathbb{Z}_p}\left(-\frac{x}{\lambda}+y\right)^l d\mu_{-q}(y)\\ &= \frac{1}{[2]_q}\sum_{l=0}^{n} S_1(n,l)(-1)^l\lambda^l E_{l,q}\left(-\frac{x}{\lambda}\right).\end{aligned} \tag{2.29}$$

When $x=0, \widehat{Bl}_{n,q}(\lambda)=\widehat{Bl}_{n,q}(0|\lambda)$ are called the q-Boole numbers of the second kind. From (2.28), we can derive the generating function of $\widehat{Bl}_{n,q}(x|\lambda)$ as follows:

$$\begin{aligned}\sum_{n=0}^{\infty}\widehat{Bl}_{n,q}(x|\lambda)\frac{t^n}{n!} &= \frac{1}{[2]_q}\int_{\mathbb{Z}_p}(1+t)^{-\lambda y+x}d\mu_{-q}(y)\\ &= \frac{(1+t)^{\lambda}}{q+(1+t)^{\lambda}}(1+t)^x.\end{aligned} \tag{2.30}$$

By replacing t by e^t-1 in (2.30), we get

$$\begin{aligned}\sum_{n=0}^{\infty}\widehat{Bl}_{n,q}(x|\lambda)\frac{(e^t-1)^n}{n!} &= \frac{e^{\lambda t}}{q+e^{\lambda t}}e^{xt}\\ &= \frac{1}{qe^{-\lambda t}+1}e^{xt}\\ &= \frac{1}{[2]_q}\sum_{m=0}^{\infty}(-1)^m\lambda^m E_{m,q}(-\frac{x}{\lambda})\frac{t^m}{m!},\end{aligned} \tag{2.31}$$

and

$$\sum_{n=0}^{\infty}\widehat{Bl}_{n,q}(x|\lambda)\frac{(e^t-1)^n}{n!} = \sum_{m=0}^{\infty}\left(\sum_{n=0}^{m}\widehat{Bl}_{n,q}(x|\lambda)S_2(m,n)\right)\frac{t^m}{m!}. \tag{2.32}$$

Therefore, by (2.31) and (2.32), we obtain the following theorem.

Theorem 2.9. *For $m\geq 0$, we have*

$$\frac{(-1)^m\lambda^m}{[2]_q}E_{m,q}(-\frac{x}{\lambda}) = \sum_{n=0}^{m}\widehat{Bl}_{n,q}(x|\lambda)S_2(m,n),$$

and

$$\widehat{Bl}_{m,q}(x|\lambda) = \frac{1}{[2]_q}\sum_{l=0}^{m} S_1(m,l)(-1)^l\lambda^l E_{l,q}\left(-\frac{x}{\lambda}\right).$$

For $k \in \mathbb{N}$, let us define the q-Boole polynomials of the second kind with order k as follows :

$$\widehat{Bl}_{n,q}^{(k)}(x|\lambda) = \frac{1}{[2]_q^k} \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} (-(\lambda x_1 + \cdots + \lambda x_k) + x)_n d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k). \tag{2.33}$$

Then we have

$$[2]_q^k \widehat{Bl}_{n,q}^{(k)}(x|\lambda) = \sum_{l=0}^{n} S_1(n,l) \lambda^l (-1)^l E_{l,q}\left(-\frac{x}{\lambda}\right).$$

From (2.33), we can derive the generating function of $\widehat{Bl}_{n,q}^{(k)}(x|\lambda)$ as follows :

$$\begin{aligned}
\sum_{n=0}^{\infty} \widehat{Bl}_{n,q}^{(k)}(x|\lambda) \frac{t^n}{n!} &= \frac{1}{[2]_q^k} \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} (1+t)^{-(\lambda x_1 + \cdots + \lambda x_k) + x} d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k) \\
&= \left(\frac{(1+t)^{\lambda}}{q + (1+t)^{\lambda}}\right)^k (1+t)^x \\
&= \left(\frac{1}{q(1+t)^{-\lambda} + 1}\right)^k (1+t)^x \\
&= \sum_{n=0}^{\infty} Bl_{n,q}^{(k)}(x|-\lambda) \frac{t^n}{n!}.
\end{aligned} \tag{2.34}$$

Thus, by (2.34), we get

$$\widehat{Bl}_{n,q}^{(k)}(x|\lambda) = Bl_{n,q}^{(k)}(x|-\lambda), (n \geq 0). \tag{2.35}$$

Indeed,

$$\begin{aligned}
(-1)^n [2]_q \frac{Bl_{n,q}(x|\lambda)}{n!} &= (-1)^n \int_{\mathbb{Z}_p} \binom{x+\lambda y}{n} d\mu_{-q}(y) \\
&= \int_{\mathbb{Z}_p} \binom{-y\lambda - x + n - 1}{n} d\mu_{-q}(y) \\
&= \sum_{m=0}^{n} \binom{n-1}{n-m} \int_{\mathbb{Z}_p} \binom{-y\lambda - x}{m} d\mu_{-q}(y) \\
&= \sum_{m=1}^{n} \frac{\binom{n-1}{m-1}}{m!} m! \int_{\mathbb{Z}_p} \binom{-y\lambda - x}{m} d\mu_{-q}(y) \\
&= [2]_q \sum_{m=1}^{n} \binom{n-1}{m-1} \frac{\widehat{Bl}_{m,q}(-x|\lambda)}{m!},
\end{aligned}$$

and

$$(-1)^n [2]_q \frac{\widehat{Bl}_{n,q}(x|\lambda)}{n!} = \sum_{m=0}^{n} \binom{n-1}{m-1} \int_{\mathbb{Z}_p} \binom{-x+y\lambda}{m} d\mu_{-q}(y)$$
$$= [2]_q \sum_{m=1}^{n} \binom{n-1}{m-1} \frac{\widehat{Bl}_{m,q}(-x|\lambda)}{m!}.$$

References

1. A. Bayad, T. Kim, *Identities involving values of Bernstein, q-Bernoulli, and q-Euler polynomials,* Russ. J. Math. Phys. **18** (2011), no. 2, 133-143.
2. D. S. Kim, T. Kim, *A note on Boole polynomials, Integral Transforms and Special Functions in 2014* DOI:10.1080/10652469.2014.891586
3. T. Kim, *Identities on the weighted q-Euler numbers and q-Bernstein polynomials,* Adv. Stud. Contemp. Math. (Kyungshang) **22** (2012), no. 1, 7-12.
4. T. Kim, *A study on the q-Euler numbers and the fermionic q-integral of the product of several type q-Bernstein polynomials on $\mathbb{Z}_p$,* Adv. Stud. Contemp. Math. (Kyungshang) **23** (2013), no. 1, 5-11.
5. T. Kim, J. Choi, Y.-H. Kim, *On extended Carlitz's type q-Euler numbers and polynomials,* Adv. Stud. Contemp. Math. (Kyungshang) **20** (2010), no. 4, 499-505.
6. T. Kim, *A note on p-adic q-integral on $\mathbb{Z}_p$ associated with q-Euler numbers,* Adv. Stud. Contemp. Math. (Kyungshang) **15** (2007), no. 2, 133-137.
7. T. Kim, *New approach to q-Euler, Genocchi numbers and their interpolation functions,* Adv. Stud. Contemp. Math. (Kyungshang) **18** (2009), no. 2, 105-112.
8. T. Kim, *Note on the q-Euler numbers of higher order,* Adv. Stud. Contemp. Math. (Kyungshang) **19** (2009), no. 1, 25-29.
9. T. Kim, B. Lee, J. Choi, Y. H. Kim, S. H. Rim, *On the q-Euler numbers and weighted q-Bernstein polynomials,* Adv. Stud. Contemp. Math. (Kyungshang) **21** (2011), no. 1, 13-18.
10. T. Kim, *The modified q-Euler numbers and polynomials,* Adv. Stud. Contemp. Math. (Kyungshang) **(**16) (2008), no. 2, 161-170.
11. S. Roman, *The umbral calculus. Pure and Applied mathematics,* Vol 111. Academic Press, Inc.[Harcourt Brace Jovanovich, publishers], New York;1984. x+193 pp. ISBN:0-12-594380-6.

1, Department of Mathematics, Sogang University, Seoul 121-742, Republic of Korea

E-mail address: dskim@kw.ac.kr

2, Department of Mathematics, Kwangwoon National University, Seoul 139-701, Republic of Korea

E-mail address: tkkim@kw.ac.kr

3, Department of Applied Mathematics, Pukyong National University, Busan 608-737, Republic of Korea

E-mail address: seo2011@pknu.ac.kr